

and gelatinization temperature, and only total protein content was slightly greater in CL1 (fig. S2). These results suggested that the CL trait has great potential for crop improvement, prompting us to explore the underlying mechanisms.

Prolonged branching time leads to increased grain number  
 During rice panicle development, a series of lateral meristems with distinct identities are sequentially generated (Fig. 1R). Inflorescence meristems first produce lateral primary branch

meristems (PBMs). Subsequently, the newly initiated meristem from the PBM either acquires the identity of secondary branch meristem (SBM) or terminates as a spikelet meristem (SM). The SBM continues to generate next-order lateral meristems that will eventually



acquire the SM identity. Finally, the SM is transformed into a floral meristem (FM) (25–27). Scanning electron microscopic (SEM) observation identified excrescent SBMs in CL1 at the early stage of panicle development (fig. 1S). In addition, the transition from BM to SM was delayed in CL1 (Fig. 1, S to U). FMs were formed in NCL1 when the young panicle of CL1 was at the SM stage (Fig. 1T), and the FMs were already fully developed in NCL1 when the FMs in the young panicle of CL1 were formed (Fig. 1U). The delay in the transition from BM to SM resulted in prolonged time for generating more SBMs and even the formation of tertiary branches in CL1 (Fig. 1E). Furthermore, an additional SM developed adjacent to the terminal spikelet (Fig. 1U). Pedicel elongation was clear in NCL1 but barely observed in CL1 (Fig. 1V). On the basis of these observations, we concluded that three developmental phenomena drive CL formation: (i) the development of more SBMs, (ii) the initiation of supernumerary SM, and (iii) shortened pedicels.

#### Cloning the causal gene through screening CL suppressors


To help us track down the genetic basis for CL, we requested three other CL germplasms (CL2, CL3, CL4) from different sources, including one (CL3, IRIS\_313-11403) from the 3000 Rice Genomes Project (3K-RG) (28). All three showed the typical CL phenotype, but they exhibited highly variable plant architecture and panicle morphology (fig. S3). We crossed them with each other and found that all F<sub>1</sub> plants derived from the different crosses retained the CL phenotype (fig. S3). By contrast, F<sub>1</sub> plants derived from crosses between CL1 and 9311, an NCL *indica* variety, or between CL1 and Zhonghua11 (ZH11), an NCL *japonica* variety, showed only a weak CL phenotype (WCL), characterized by clustered growth of two grains at the terminus of the panicle branches (figs. S4 and S5). In addition, in the crossed CL1 × ZH11 F<sub>2</sub> population, the segregation ratios (NCL:WCL:CL) were 65:131:45 (fig. S5), close to 1:2:1 ( $\chi^2 = 0.0951664 < 5.99$  at  $P = 0.05$ ). These results suggested that the distinctive phenotype of these CL plants is determined by the same semidominant locus. However, our intense efforts using either map-based cloning or bulked-segregant analysis (BSA) by generating large populations or different crosses could only locate the locus to the ~22.85- to 23.85-Mb (megabase) region on chromosome 6 (fig. S6), like previously reported attempts (21–24). One possibility is that there exist complex chromosome structural variations around the region.

The failure of these routine cloning methods prompted us to design a strategic cloning roadmap by generating large-scale mutagenesis of CL1 for identifying the causal gene (Fig. 2A). From a population of 10,000 inde-


pendent mutant lines (M<sub>1</sub>) using sodium azide as the mutagen, examination of the phenotypes of the M<sub>2</sub> plants (16 for each line) led to the identification of two mutant lines that reverted to a wild-type NCL phenotype, which we designated *cll-1* and *cll-2* (Fig. 2B and fig. S7). By comparing the NCL and CL plants pooled from the F<sub>2</sub> population of the *cll* × CL1 backcrosses, we found that, similar to NCL1, both *cll-1* and *cll-2* had significantly decreased grain number compared to CL1, owing to their smaller numbers of secondary branches (Fig. 2, D to F). Both mutants also showed identical plant architecture, panicle number, panicle length, and 1000-grain weight to those of NCL1 and CL1 (figs. S8 and S9 and Fig. 2G). In addition, the grain yield per plant of the two *cll* mutants decreased by 16.4% to 27.7% compared to CL1 in Sanya (Fig. 2H). Because CL is controlled by a single semidominant



**I** **J**









**L** **M**



no detectable differences in grain size and plant height among CL1, *cli-1*, *cli-2*, and NCL1 (Fig. 3, C to E, and fig. S17). Because CL is closely associated with shortened pedicels, we first compared the expression pattern of *BRD3* in pedicels and spikelets of NCL1 and CL1. Notably, *BRD3* was up-regulated in pedicels, but

not spikelets, of CL1 (Fig. 3F). To confirm this result, we also measured BR content in these tissues of both lines. Consistent with the expression pattern of *BRD3*, the castasterone level was only ~20% lower in CL1 spikelets, yet ~70% lower in CL1 pedicels, compared to NCL1 spikelets and pedicels (Fig. 3G).


We further conducted RNA in situ hybridization to determine the spatiotemporal expression pattern of *BRD3* during panicle development. During the development of NCL1 inflorescences, *BRD3* expression was very weak and not specific to any tissues (Fig. 3, H and I). By contrast, we detected specific signal with



top-down development in SBM, but not PBM, of CL1 panicles (Fig. 3, H and I). At the stage of SM or FM development, *BRD3* expression was specifically detected at the base of spikelets, corresponding to the pedicel positions, in CL1 but not NCL1 (Fig. 3, J to L). As a negative control, no signal was detected when *BRD3* was used as the sense probe (fig. S18). The strong relationship between *BRD3* expression pattern and plant phenotype suggests that spatial enhancement of *BRD3* expression gives rise to the enhanced grain number, with no impact on grain size.

Spatial expression of GSK2 produces CL

Because spatial up-regulation of *BRD3* in CL1 suppresses BR biosynthesis and BRs function through the BR signaling pathway, spatial suppression of BR signaling could also enable



**Fig. 4. Spatially specific expression of GSK2 produces CL.** (A) Panicles of ZH11 and the transgenic plants. Scale bars: 2 cm. (B) Different cluster types in FLAG-aGSK2 plants. Scale bar: 1 cm. (C) Detection of the fusion proteins in the transgenic plants by immunoblotting analyses. ACTIN was detected as control. M (kD), marker size indicated in kilodaltons. (D) Percentages of grain numbers belonging to different cluster types in the three FLAG-aGSK2 lines, with 15 panicles analyzed for each line. (E) Comparison of grain size in different GSK2-related plants. Scale bar: 5 mm.









(F and G) Detection of the fusion proteins in spikelets (S) and pedicels (P) of the plants by immunoblotting. Asterisk (\*) represents a nonspecific band indicating equal protein loading. Arrowheads indicate target protein signals. (H to K) Immunofluorescence analysis of the fusion proteins during inflorescence development in FLAG-aGSK2 and GFP-aGSK2. Ped, pedicel. The expression in the presumed pedicel positions were circled. Scale bars: 100  $\mu$ m. (L and M) Comparison of the panicles and primary branches of CL1, FLAG-aGSK2, and the double mutant. Scale bars: 2 cm (L) and 1 cm (M).

OsmADS1-FLAG fusion protein (fig. S29). These results strongly suggested that phosphorylation of OsmADS1 by GSK2 enhances the stability of the OsmADS1 protein.

OsmADS1 accumulates in pedicels of CLs. Given that GSK2 targets and promotes OsmADS1, the OsmADS1 protein should also specifically accumulate in CL plants in the same way that

FLAG-aGSK2 protein or BRD3 transcripts do. To test this notion, we analyzed OsmADS1 protein levels in spikelets and pedicels of FLAG-aGSK2 and GFP-aGSK2 lines. Compared to wild






type, in spikelets, OsMADS1 was more abundant in *GFP-aGSK2*, but of similar abundance in *FLAG-aGSK2* (Fig. 5G). By contrast, in pedicels, OsMADS1 strongly accumulated in *FLAG-aGSK2*, but not in *GFP-aGSK2* (Fig. 5H). These data clearly demonstrated that the distribution patterns of OsMADS1 and GSK2 are highly similar in either *GFP-aGSK2* or *FLAG-aGSK2* (Figs. 4, F and G, and 5, G and H). In addition, the distribution patterns are also consistent with the plant phenotypes: *GFP-aGSK2* has smaller grains, whereas *FLAG-aGSK2* has shortened pedicels and normal grain size (Fig. 4E).



We then extended the analysis to CL1 and NCL1. Consistently, OsMADS1 levels were higher in pedicels, but not spikelets, of CL1 compared to NCL1 (Fig. 5, I and J). These patterns were similar to those observed in *FLAG-aGSK2*. Taken together, these results demonstrated that OsMADS1 acts downstream of GSK2 to mediate BR regulation of CL development.








Knockdown of **O. MAD 4** abolishes CL phenotype

Similar to GSK2, OsMADS1 has been characterized as a negative regulator of grain size (35, 36). Furthermore, OsMADS1 has been implicated in regulating meristem identity and potentially enhancing spikelet number (37-39). Examination of *OsMADS1*-overexpressing plants (



**J**





number with 81.5 and 39.7% more grains per panicle compared to R498 and CL1, respectively (Fig. 7, E and F). In addition, CL5 possessed other advantageous traits such as a thick culm, heavier grains (fig. S35), and high yield (Fig. 7G), illustrating the promising potential of combining *CL* and *Gn1a* for enhancing rice yield.

BR alteration in clustered growth in other species

Although we demonstrated that BR depletion controls CL in rice (Fig. 7H), we wanted to know whether this represent a general mechanism underlying clustered growth across plant species. Cluster peppers (*Capsicum annuum* L. var. *fasciculatum*) display multiple flowers clustered on a single flower bud formation node, whereas common peppers (*Capsicum annuum* L.) have only one flower attached to the node. We measured the BR content in pedicels of Tianyu 2 (TY2), a common pepper, and TY2<sup>CL</sup>, a cluster pepper showing clustered flowers and shortened pedicels while closely resembling TY2 in vegetative and reproductive morphology (Fig. 7, I and J). Like CL1 and NCL1, the castasterone level was lower in TY2<sup>CL</sup> pedicels compared to TY2 (Fig. 7K). Conversely, the 6-deoxocastasterone level, the precursor of castasterone, was higher in TY2<sup>CL</sup> pedicels (Fig. 7K), indicating a possible loss of function in the BR synthase responsible for converting 6-deoxocastasterone to castasterone.

We also measured the BR content in the pedicels of the vine *Rosa chinensis* (*Rosa* “Parkdirektor Riggers”) and the vine *Rosa* sp. (*Rosa* “Angela”), which displays the clustered flowers and shortened pedicels (Fig. 7, L and M). Like TY2 and TY2<sup>CL</sup>, the castasterone level in *Rosa* “Angela” pedicels was lower compared to *Rosa* “Parkdirektor Riggers,” whereas the 6-deoxocastasterone level exhibited the opposite trend (Fig. 7N). These consistent findings indicate that BR distribution may play a general role in regulating the clustered growth and inflorescence structure.

## Discussion

We unraveled the genetic basis underlying the historic CL germplasm, shedding light on the development of CL and its association with enhanced grain number, a crucial determinant of yield. We demonstrated that defective BR catabolism or signaling contributes to spikelet clustering and increased grain number, mediated by the specific spatial expression of *BRD3* transcripts, GSK2 protein, OsMADS1 protein, and *RCN2* transcripts. During CL's panicle development, the activation of *BRD3* in SBM triggers BR degradation, which enhances GSK2 stability, promotes OsMADS1 accumulation, and subsequently increases *RCN2* expression. As a result, the identity of SM is suppressed, leading to a delayed transition from BM to SM, ultimately yielding more SBMs and additional SMs formed adjacent to the ter-

minal spikelet (Fig. 7H). Overall, our study exemplifies the successful utilization of natural rice germplasm resources harboring beneficial alleles to breed superior rice strains, from gene cloning to the dissection of the underlying molecular mechanisms.

KNOX transcription factors play a crucial role in the establishment and maintenance of the shoot apical meristem (SAM) (48). In *Arabidopsis*, BREVIPEDICELLUS (BP), a KNOX transcription factor, has been shown to regulate pedicel development (48). In rice, ORYZA SATIVA HOMEBOX1 (OSH1), the closest ortholog of BP, targets and activates BR catabolic genes, including *BRD3*, to maintain SAM activity (49). Therefore, the degradation of BRs should also promote the activity of SBM and SM. In our study, we observed an increase in grain number accompanied by shortened pedicels, and the expression of *BRD3* was detected sequentially in both the meristem cells and pedicel cells, suggesting a potential derivation of pedicels from meristem cells. Our findings imply that large structural variations preceding the *BRD3* gene may be responsible for the spatially specific increase in *BRD3* expression observed in CL. Although the detailed mechanism requires further exploration, one possibility is that these structural variations enhance chromatin accessibility, allowing the entry of transcription factors, such as OSH1, to promote *BRD3* expression.

Constitutive activation of *BRD3* in a transferred DNA (T-DNA) insertion mutant results in severe developmental defects, whereas moderately increased expression of *BRD3* in the heterozygote leads to higher grain number but smaller grains (29). Therefore, the practical application of *CL* in crop improvement can be attributed to two main factors related to *BRD3* expression: spatially specific expression and optimal expression. The resulting spatial distribution of BRs provides beneficial effects while preventing the reduction in grain size typically observed in BR-defective mutants, showcasing its practical applicability in crop improvement. Together with our previous research on the distribution of cytokinins and auxins (50, 51), this work highlights the potential of optimizing hormone distributions to achieve desirable traits while mitigating negative associations in crop improvement efforts. Improving grain yield is intricately linked to expanding source supply, influenced by various factors such as source capacity, growth conditions (light, water, temperature, nutrients), and fertilizers. Although introduction of CL into several different modern cultivars all resulted in yield improvement, we observed distinct effects in different locations and genetic backgrounds. Considering that CL is influenced by BR content and each rice may exhibit variable BR levels, it is crucial to extensively test the effectiveness of CL in diverse contexts.

## Materials and methods summary

Plant growth, suppressor screening, and segregating populations

For field plot analysis, each material was planted with a row interspace of 20 cm and plant interspace of 17.1 cm, with ~33 plants per square meter. At least four plots were planted as biological replicates. The border plants were removed from each plot to avoid margin effects during measurements. For experiments requiring seedling analysis, plants were grown in a growth chamber at 30°C with 10 hours light for day and 28°C with 14 hours dark for night. Half-strength Murashige and Skoog (½-MS) solution was provided as the nutrient source. For suppressor screening, dry CL1 seeds were incubated in sodium azide (1 mM in phosphate buffer pH 3.0) for 6 hours. After a thorough wash with tap water, the seeds were germinated and planted. A total of 10,000 individual lines were harvested, and 16 plants of each line were grown for mutant screening. Four segregating populations were generated by crossing CL1 with ZH11, 9311, *cll-1*, and *cll-2*, respectively. For each population, pooled DNAs from NCL plants and CL plants were prepared for BSA. Numbers of individual plants used for DNA pools from each population were as follows: 36 NCL and 34 CL from CL1 × ZH11, 24 NCL and 31 CL from CL1 × 9311, 31 NCL and 33 CL from CL1 × *cll-1*, and 20 NCL and 19 CL from CL1 × *cll-2*.

RNA in situ hybridization

Specific fragments of *BRD3* and *RCN2* were amplified by PCR with primers listed in table S1. The fragments were subsequently inserted into the pEASY-Blunt Simple Cloning Vector (TransGen) for in vitro RNA transcription. Sense and antisense RNA probes were produced using T7 transcriptase and labeled with digoxigenin (Roche). Young panicle tissues from NCL1 and CL1 were collected and fixed in FAA (45% ethanol, 6% glacial acetic acid, and 5% formaldehyde) at 4°C overnight in a vacuum, dehydrated, and embedded in Paraplast Plus (Sigma-Aldrich, St. Louis). Tissue sections were prepared using a microtome (RM2235; Leica, Wetzlar) and then affixed to Poly-Prep slides (Sigma-Aldrich, St. Louis).

In vitro phosphorylation and mass spectrum analysis

One microgram of GST-GSK2 and 2 µg of MBP-OsMADS1 or MBP-OsMADS1<sup>TA</sup> were incubated in the kinase buffer [10 mM Tris-HCl (pH 7.4), 50 mM NaCl, 6 mM MgCl<sub>2</sub>, 0.5 mM dithiothreitol] with or without ATP and CIP (NEB) at 30°C for 2 hours. Phos-tag<sup>TM</sup> SDS-polyacrylamide gel electrophoresis (PAGE) was used to separate the phosphorylation protein following the instruction of the product Phos-tag<sup>TM</sup> Acrylamide (Wako). To identify the phosphorylation sites, 4 µg of recombinant

MBP-OsMADS1 or MBP-OsMADS1<sup>TA</sup> proteins were phosphorylated by 2 µg of GST-GSK2 for 1 hour in vitro. Phos-tag<sup>TM</sup> SDS-PAGE was used to separate the phosphorylation protein. The target protein bands were cut after one-step blue staining (Biotium). The LC-MS/MS detection and analysis were conducted by the Shanghai Luming Biological Technology Co., Ltd. (Shanghai, China).

#### Immunofluorescence detection

An 8-µm-thick section was cut from paraffin-embedded young panicle tissues for detection. The slides were deparaffinized, rehydrated, and subjected to epitope retrieval by boiling in 1× All-purpose Powerful Antigen Retrieval Solution (Beyotime) for 20 min at 95°C, and set to cool for 30 to 40 min. The slides were washed two times in 1× Immunol Staining Wash Buffer (Beyotime) for 10 min each, blocked in Immunol Staining Blocking Buffer (Beyotime) for 60 min at room temperature, and then rinsed two times in 1× Immunol Staining Wash Buffer (Beyotime) for 10 min each. The primary antibody for OsMADS1 (1:50; Abclonal, Cat#A20328), or for FLAG-aGSK2 (anti-FLAG, 1:50; Sigma, Cat#F1804), or for GFP-aGSK2 (anti-GFP, 1:50; Abmart, Cat#M2004I), was used for incubation overnight at 4°C. The specimen was rinsed three times in 1× Immunol Staining Wash Buffer (Beyotime) for 5 min each, and then incubated in the Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor<sup>TM</sup> Plus 555 (1:400; Invitrogen, Cat#A32732) or Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488 (1:400; Invitrogen, Cat#A32723) at room temperature for 60 min in dark. The slides were further rinsed three times in 1× Immunol Staining Wash Buffer (Beyotime) for 5 min each, and then covered using coverslips filled with Antifade Mounting Medium with 4',6-diamidino-2-phenylindole (DAPI; Beyotime). Fluorescence was captured with a confocal laser scanning microscope (LSM 980; Zeiss, Oberkochen).

#### ChIP-seq and RNA-seq

About 1 g of young panicles at the 5-mm stage of *OsMADS1-OE* were used for ChIP-seq. Tissue fixation, nuclei extraction, and chromatin immunoprecipitation were performed using anti-OsMADS1 (1:100; Abclonal, Cat#A20328) antibody. The ChIP-seq DNA libraries were sequenced using the Illumina HiSeqTM2000 platform. After quality control, BWA software was used to align the clean reads against the Nipponbare reference genome (IRGSP1.0). MACS software was used for peak calling on a genome-wide basis, and the threshold for screening significant peak was  $q$ -value < 0.05. Significant peaks were assigned to the nearest gene. For RNA-seq, about 1 g of young panicles

at the 5-mm stage of *OsMADS1-OE* and ZH11 were collected, and total RNA was extracted from each of three biological replicates using TRIzol reagent (Invitrogen). Sequencing libraries were generated using NEBNext Ultra RNA Library Prep Kit for Illumina (NEB), following the manufacturer's recommendations, and sequenced on Illumina NovaSeq platform. After cleaning up raw sequence reads, the clean reads were mapped to the Nipponbare reference genome (IRGSP1.0). The differentially expressed genes were analyzed using the edgeR package. Genes with a false discovery rate (FDR) < 0.05 and log<sub>2</sub>(fold change) > 1 were assigned as differentially expressed.

#### REFERENCES AND NOTES

1. T. Sasaki, B. Burr, International Rice Genome Sequencing Project: The effort to completely sequence the rice genome. *Curr. Opin. Plant Biol.* **3**, 138–142 (2000). doi: [10.1016/S1369-5266\(99\)00047-3](https://doi.org/10.1016/S1369-5266(99)00047-3); pmid: [10712951](https://pubmed.ncbi.nlm.nih.gov/10712951/)
2. X. Huang et al., Natural variation at the *DEPI* locus enhances grain yield in rice. *Nat. Genet.* **41**, 494–497 (2009). doi: [10.1038/ng.352](https://doi.org/10.1038/ng.352); pmid: [19305410](https://pubmed.ncbi.nlm.nih.gov/19305410/)
3. M. Ashikari et al., Cytokinin oxidase regulates rice grain production. *Science* **309**, 741–745 (2005). doi: [10.1126/science.1113373](https://doi.org/10.1126/science.1113373); pmid: [15976269](https://pubmed.ncbi.nlm.nih.gov/15976269/)
4. E. J. Kim, E. Russinova, Brassinosteroid signalling. *Curr. Biol.* **30**, R294–R298 (2020). doi: [10.1016/j.cub.2020.02.011](https://doi.org/10.1016/j.cub.2020.02.011); pmid: [32259497](https://pubmed.ncbi.nlm.nih.gov/32259497/)
5. H. Tong, C. Chu, Functional specificities of brassinosteroid and potential utilization for crop improvement. *Trends Plant Sci.* **23**, 1016–1028 (2018). doi: [10.1016/j.tplants.2018.08.007](https://doi.org/10.1016/j.tplants.2018.08.007); pmid: [30220494](https://pubmed.ncbi.nlm.nih.gov/30220494/)
6. T. M. Nolan, N. Vukašinović, D. Liu, E. Russinova, Y. Yin, Brassinosteroids: Multidimensional regulators of plant growth, development, and stress responses. *Plant Cell* **32**, 295–318 (2020). doi: [10.1105/tpc.19.00335](https://doi.org/10.1105/tpc.19.00335); pmid: [31776234](https://pubmed.ncbi.nlm.nih.gov/31776234/)
7. H. Tong et al., DWARF AND LOW-TILLERING acts as a direct downstream target of a GSK3/SHAGGY-like kinase to mediate brassinosteroid responses in rice. *Plant Cell* **24**, 2562–2577 (2012). doi: [10.1105/tpc.112.097394](https://doi.org/10.1105/tpc.112.097394); pmid: [22685166](https://pubmed.ncbi.nlm.nih.gov/22685166/)
8. J. Li, K. H. Nam, Regulation of brassinosteroid signaling by a GSK3/SHAGGY-like kinase. *Science* **295**, 1299–1301 (2002). doi: [10.1126/science.1065769](https://doi.org/10.1126/science.1065769); pmid: [11847343](https://pubmed.ncbi.nlm.nih.gov/11847343/)
9. J. H. Youn, T. W. Kim, Functional insights of plant GSK3-like kinases: Multi-taskers in diverse cellular signal transduction pathways. *Mol. Plant* **8**, 552–565 (2015). doi: [10.1016/j.molp.2014.12.006](https://doi.org/10.1016/j.molp.2014.12.006); pmid: [25655825](https://pubmed.ncbi.nlm.nih.gov/25655825/)
10. Z. Hong et al., A rice brassinosteroid-deficient mutant, *ebisu dwarf* (*d2*), is caused by a loss of function of a new member of cytochrome P450. *Plant Cell* **15**, 2900–2910 (2003). doi: [10.1105/tpc.014712](https://doi.org/10.1105/tpc.014712); pmid: [14615594](https://pubmed.ncbi.nlm.nih.gov/14615594/)
11. S. Tanabe et al., A novel cytochrome P450 is implicated in brassinosteroid biosynthesis via the characterization of a rice dwarf mutant, *dwarf11*, with reduced seed length. *Plant Cell* **17**, 776–790 (2005). doi: [10.1105/tpc.104.024950](https://doi.org/10.1105/tpc.104.024950); pmid: [15705958](https://pubmed.ncbi.nlm.nih.gov/15705958/)
12. C. Yamamuro et al., Loss of function of a rice brassinosteroid insensitive1 homolog prevents internode elongation and bending of the lamina joint. *Plant Cell* **12**, 1591–1606 (2000). doi: [10.1105/tpc.12.9.1591](https://doi.org/10.1105/tpc.12.9.1591); pmid: [11006334](https://pubmed.ncbi.nlm.nih.gov/11006334/)
13. L. Song et al., Reducing brassinosteroid signalling enhances grain yield in semi-dwarf wheat. *Nature* **617**, 118–124 (2023). doi: [10.1038/s41586-023-06023-6](https://doi.org/10.1038/s41586-023-06023-6); pmid: [37100915](https://pubmed.ncbi.nlm.nih.gov/37100915/)
14. J. Tian et al., Teosinte ligule allele narrows plant architecture and enhances high-density maize yields. *Science* **365**, 658–664 (2019). doi: [10.1126/science.aax5482](https://doi.org/10.1126/science.aax5482); pmid: [31416957](https://pubmed.ncbi.nlm.nih.gov/31416957/)
15. T. Sakamoto et al., Erect leaves caused by brassinosteroid deficiency increase biomass production and grain yield in rice. *Nat. Biotechnol.* **24**, 105–109 (2006). doi: [10.1038/nbt1173](https://doi.org/10.1038/nbt1173); pmid: [16369540](https://pubmed.ncbi.nlm.nih.gov/16369540/)
16. K. Ramiah, S. Jobitharaj, S. D. Mudaliar, Inheritance of characters in rice, Part IV. *Mem. Dept. Agric. India Bot. Ser.* **18**, 229–259 (1931).
17. N. E. Jodon, Inheritance and linkage relationships of a chlorophyll mutation in rice. *Agron. J.* **32**, 342–346 (1940). doi: [10.2134/agronj1940.00021962003200050004x](https://doi.org/10.2134/agronj1940.00021962003200050004x)

18. N. E. Jodon, Inheritance of some of the more striking characters in rice. *J. Hered.* **48**, 181–192 (1957). doi: [10.1093/oxfordjournals.jhered.a106718](https://doi.org/10.1093/oxfordjournals.jhered.a106718)
19. S. Nagao, M. E. Takahashi, Genetical studies on rice plant, XXVII. Trial construction of twelve linkage groups in Japanese rice. *J. Fac. Agric. Hokkaido Univ.* **53**, 72–130 (1963).
20. H. Chen, G. Liu, X. Zhu, S. Min, Observation and genetically analysis on character of clustered spikelets in rice. *J. Nanjing Agric. Univ.* **25**, 116–118 (2002).
21. C. Tian et al., Genetic analysis and preliminary gene mapping of rice clustered spikelet mutant. *Mol. Plant Breed.* **8**, 29–34 (2010).
22. Y. Zhang et al., Development of NILs with *Cl*-gene of rice restorer and evaluation on the near-isogenic level. *Acta Agron. Sin.* **32**, 397–401 (2006).
23. L. Y. Zheng et al., Morphology and mapping analysis of rice (*Oryza sativa* L.) clustered spikelets (*Cl*) mutant. *Chin. Sci. Bull.* **48**, 559–562 (2003). doi: [10.1360/03tb9119](https://doi.org/10.1360/03tb9119)
24. Y. Zheng et al., Fine mapping and candidate gene analysis of clustered spikelet gene *OsCl-6* in rice (*Oryza sativa*). *J. Agric. Biotechnol.* **26**, 1116–1123 (2018).
25. M. Nakagawa, K. Shimamoto, J. Kyoizuka, Overexpression of *RCN1* and *RCN2*, rice *TERMINAL FLOWER 1/CENTRORADIALIS* homologs, confers delay of phase transition and altered panicle morphology in rice. *Plant J.* **29**, 743–750 (2002). doi: [10.1046/j.1365-3113.2002.01255.x](https://doi.org/10.1046/j.1365-3113.2002.01255.x); pmid: [12148532](https://pubmed.ncbi.nlm.nih.gov/12148532/)
26. P. Prusinkiewicz, Y. Erasmus, B. Lane, L. D. Harder, E. Coen, Evolution and development of inflorescence architectures. *Science* **316**, 1452–1456 (2007). doi: [10.1126/science.1140429](https://doi.org/10.1126/science.1140429); pmid: [17525303](https://pubmed.ncbi.nlm.nih.gov/17525303/)
27. P. McSteen, D. Laudencia-Chingcuanco, J. Colasanti, A floret by any other name: Control of meristem identity in maize. *Trends Plant Sci.* **5**, 61–66 (2000). doi: [10.1016/S1360-1385\(99\)01541-1](https://doi.org/10.1016/S1360-1385(99)01541-1); pmid: [10664615](https://pubmed.ncbi.nlm.nih.gov/10664615/)
28. W. Wang et al., Genomic variation in 3,010 diverse accessions of Asian cultivated rice. *Nature* **557**, 43–49 (2018). doi: [10.1038/s41586-018-0063-9](https://doi.org/10.1038/s41586-018-0063-9); pmid: [29695866](https://pubmed.ncbi.nlm.nih.gov/29695866/)
29. W. Qian et al., Novel rice mutants overexpressing the brassinosteroid catabolic gene *CYP73A4A*. *Plant Mol. Biol.* **93**, 197–208 (2017). doi: [10.1007/s11033-016-0558-4](https://doi.org/10.1007/s11033-016-0558-4); pmid: [27815670](https://pubmed.ncbi.nlm.nih.gov/27815670/)
30. T. Sakamoto et al., Rice CYP73A4s function as multisubstrate and multifunctional enzymes in brassinosteroid catabolism. *Plant J.* **67**, 1–12 (2011). doi: [10.1111/j.1365-3113.2011.04567.x](https://doi.org/10.1111/j.1365-3113.2011.04567.x); pmid: [21418356](https://pubmed.ncbi.nlm.nih.gov/21418356/)
31. H. A. Agrama et al., Genetic assessment of a mini-core subset developed from the USDA rice genebank. *Crop Sci.* **49**, 1336–1346 (2009). doi: [10.2135/cropsci2008.06.0551](https://doi.org/10.2135/cropsci2008.06.0551)
32. D. Liu et al., Diversification of plant agronomic traits by genome editing of brassinosteroid signaling family genes in rice. *Plant Physiol.* **187**, 2563–2576 (2021). doi: [10.1093/plphys/kiab394](https://doi.org/10.1093/plphys/kiab394); pmid: [34618079](https://pubmed.ncbi.nlm.nih.gov/34618079/)
33. H. Tong et al., Brassinosteroid regulates cell elongation by modulating gibberellin metabolism in rice. *Plant Cell* **26**, 4376–4393 (2014). doi: [10.1105/tpc.114.132092](https://doi.org/10.1105/tpc.114.132092); pmid: [25371548](https://pubmed.ncbi.nlm.nih.gov/25371548/)
34. L. Wang et al., Ectopic expression of *OsMADS1* caused dwarfism and spikelet alteration in rice. *Plant Growth Regul.* **81**, 433–442 (2017). doi: [10.1007/s10725-016-0220-9](https://doi.org/10.1007/s10725-016-0220-9)
35. Q. Liu et al., G-protein β subunits determine grain size through interaction with MADS-domain transcription factors in rice. *Nat. Commun.* **9**, 852 (2018). doi: [10.1038/s41467-018-03047-9](https://doi.org/10.1038/s41467-018-03047-9); pmid: [29487282](https://pubmed.ncbi.nlm.nih.gov/29487282/)
36. J. Yu et al., Alternative splicing of *OsLG3b* controls grain length and yield in japonica rice. *Plant Biotechnol. J.* **16**, 1667–1678 (2018). doi: [10.1111/pbi.12903](https://doi.org/10.1111/pbi.12903); pmid: [29479793](https://pubmed.ncbi.nlm.nih.gov/29479793/)
37. J. S. Jeon et al., *leafy hull sterile1* is a homeotic mutation in a rice MADS box gene affecting rice flower development. *Plant Cell* **12**, 871–884 (2000). doi: [10.1105/tpc.12.6.871](https://doi.org/10.1105/tpc.12.6.871); pmid: [10852934](https://pubmed.ncbi.nlm.nih.gov/10852934/)
38. K. Prasad, S. Parameswaran, U. Vijayraghavan, *OsMADS1*, a rice MADS-box factor, controls differentiation of specific cell types in the lemma and palea and is an early-acting regulator of inner floral organs. *Plant J.* **43**, 915–928 (2005). doi: [10.1111/j.1365-3113.2005.02504.x](https://doi.org/10.1111/j.1365-3113.2005.02504.x); pmid: [16146529](https://pubmed.ncbi.nlm.nih.gov/16146529/)
39. D. Ren, Y. Li, G. He, Q. Qian, Multifloret spikelet improves rice yield. *New Phytol.* **225**, 2301–2306 (2020). doi: [10.1111/nph.16303](https://doi.org/10.1111/nph.16303); pmid: [31677165](https://pubmed.ncbi.nlm.nih.gov/31677165/)
40. S. T. Malcomber, E. A. Kellogg, Heterogeneous expression patterns and separate roles of the *SEPALLATA* gene *LEAFY HULL STERILE1* in grasses. *Plant Cell* **16**, 1692–1706 (2004). doi: [10.1105/tpc.021576](https://doi.org/10.1105/tpc.021576); pmid: [15208396](https://pubmed.ncbi.nlm.nih.gov/15208396/)
41. K. Prasad, P. Sriram, C. S. Kumar, K. Kushalappa, U. Vijayraghavan, Ectopic expression of rice *OsMADS1* reveals

- a role in specifying the lemma and palea, grass floral organs analogous to sepals. *Dev. Genes Evol.* **211**, 281–290 (2001). doi: [10.1007/s004270100153](https://doi.org/10.1007/s004270100153); pmid: [11466523](https://pubmed.ncbi.nlm.nih.gov/11466523/)
42. I. Khanday, S. R. Yadav, U. Vijayraghavan, Rice *LHSL/OsMADS1* controls floret meristem specification by coordinated regulation of transcription factors and hormone signaling pathways. *Plant Physiol.* **161**, 1970–1983 (2013). doi: [10.1104/pp.112.212423](https://doi.org/10.1104/pp.112.212423); pmid: [23449645](https://pubmed.ncbi.nlm.nih.gov/23449645/)
43. J. T. Robinson *et al.*, Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011). doi: [10.1038/nbt.1754](https://doi.org/10.1038/nbt.1754); pmid: [21221095](https://pubmed.ncbi.nlm.nih.gov/21221095/)
44. W. Zhu *et al.*, Rice *SEPALLATA* genes *OsMADS5* and *OsMADS34* cooperate to limit inflorescence branching by repressing the *TERMINAL FLOWER1*-like gene *RCN4*. *New Phytol.* **233**, 1682–1700 (2022). doi: [10.1111/nph.17855](https://doi.org/10.1111/nph.17855); pmid: [34767634](https://pubmed.ncbi.nlm.nih.gov/34767634/)
45. S. Zhang *et al.*, *TFL1/CEN*-like genes control intercalary meristem activity and phase transition in rice. *Plant Sci.* **168**, 1393–1408 (2005). doi: [10.1016/j.plantsci.2004.10.022](https://doi.org/10.1016/j.plantsci.2004.10.022)
46. X. Yang *et al.*, miR156f integrates panicle architecture through genetic modulation of branch number and pedicel length pathways. *Rice (N. Y.)* **12**, 40 (2019). doi: [10.1186/s12284-019-0299-5](https://doi.org/10.1186/s12284-019-0299-5); pmid: [31147794](https://pubmed.ncbi.nlm.nih.gov/31147794/)
47. B. Tu *et al.*, Loss of *Gn1a/OsCKX2* confers heavy-panicle rice with excellent lodging resistance. *J. Integr. Plant Biol.* **64**, 23–38 (2022). doi: [10.1111/jipb.13185](https://doi.org/10.1111/jipb.13185); pmid: [34783157](https://pubmed.ncbi.nlm.nih.gov/34783157/)
48. S. J. Douglas, C. D. Riggs, Pedicel development in *Arabidopsis thaliana*: Contribution of vascular positioning and the role of the *BREVIPEDICELLUS* and *ERECTA* genes. *Dev. Biol.* **284**, 451–463 (2005). doi: [10.1016/j.ydbio.2005.06.011](https://doi.org/10.1016/j.ydbio.2005.06.011); pmid: [16038894](https://pubmed.ncbi.nlm.nih.gov/16038894/)
49. K. Tsuda, N. Kurata, H. Ohyanagi, S. Hake, Genome-wide study of *KNOX* regulatory network reveals brassinosteroid catabolic genes important for shoot meristem function in rice. *Plant Cell* **26**, 3488–3500 (2014). doi: [10.1105/tpc.114.129122](https://doi.org/10.1105/tpc.114.129122); pmid: [25194027](https://pubmed.ncbi.nlm.nih.gov/25194027/)
50. L. Liu *et al.*, Activation of *Big Grain1* significantly improves grain size by regulating auxin transport in rice. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 11102–11107 (2015). doi: [10.1073/pnas.1512748112](https://doi.org/10.1073/pnas.1512748112); pmid: [26283354](https://pubmed.ncbi.nlm.nih.gov/26283354/)
51. W. Yin *et al.*, ARGONAUTE2 enhances grain length and salt tolerance by activating *BIG GRAIN3* to modulate cytokinin distribution in rice. *Plant Cell* **32**, 2292–2306 (2020). doi: [10.1105/tpc.19.00542](https://doi.org/10.1105/tpc.19.00542); pmid: [32409321](https://pubmed.ncbi.nlm.nih.gov/32409321/)
52. X. Zhang, Nanopore sequencing, RNA-seq and ChIP-seq data. *Zenodo* (2024); doi: [10.5281/zenodo.10574305](https://doi.org/10.5281/zenodo.10574305)
53. X. Zhang, BSA data. *Zenodo* (2024); doi: [10.5281/zenodo.10574366](https://doi.org/10.5281/zenodo.10574366)
54. X. Zhang, Resequencing data of CL1. *Zenodo* (2024); doi: [10.5281/zenodo.10574479](https://doi.org/10.5281/zenodo.10574479)

#### ACKNOWLEDGMENTS

We thank Q. Li (Yangzhou University) for grain quality evaluation, Y. Li (Southwest University) for providing *OsMADS1-OE* lines, C. Sun (Sichuan Agricultural University) for sequence resemblance, G. Jiang (IGDB-CAS) and W. Wang (ICS-CAAS) for providing CL plants, H. Wu (IVF-CAAS) for providing peppers, C. Tian (IVF-CAAS) for suggestion of rose, and B. Cai (Wuhan Greensword Creation Technology Co. Ltd.) for BR quantification. **Funding:** This work was supported by National Natural Science Foundation (U21A20208, 32272141), Hainan Seed Industry Laboratory

(B23CJ0208), Innovation Program of Chinese Academy of Agricultural Sciences, and “5511” Collaborative Innovation Project (XTCXGC2021001). **Author contributions:** H.T. conceived and designed the study. X.Z., W.M., and D.L. performed experiments. Y.Y., Z.C., X.M., W.Y., M.N., N.D., and J.L. and Z.L. provided assistance. D.P., W.S., Y.Q.L., and M.Z. conducted field tests and breeding. X.Z., W.M., and H.T. analyzed the data and wrote the manuscript. H.T., Q.Q., M.Z., and C.C. cosupervised the study. **Competing interests:** H.T., X.Z., D.L., W.M., W.Y., M.N., N.D., J.L., and Y.Y. are inventors on a granted patent in China related to this work (no. ZL202210239578.7). The other authors declare that they have no competing interests. **Data and materials availability:** All sequencing data generated in this paper have been deposited and made available at Zenodo (52–54). All other data are available in the main text or the supplementary materials. **License information:** Copyright © 2024 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.sciencemag.org/about/science-licenses-journal-article-reuse>

#### SUPPLEMENTARY MATERIALS

[science.org/doi/10.1126/science.adk8838](https://science.org/doi/10.1126/science.adk8838)

Materials and Methods

Figs. S1 to S35

Tables S1 and S2

References (55–73)

MDAR Reproducibility Checklist

Submitted 15 September 2023; accepted 29 January 2024  
10.1126/science.adk8838